

February 1, 2008

A quark model analysis of the transversity distribution: Next to leading order evolution[†]

Sergio Scopetta and Vicente Vento^(a)

*Departament de Física Teòrica
Universitat de València
46100 Burjassot (València), Spain*

and

*(a) Institut de Física Corpuscular
Consejo Superior de Investigaciones Científicas*

Abstract

The feasibility of measuring chiral-odd parton distribution functions in polarized Drell-Yan and semi-inclusive experiments has renewed theoretical interest in their study. Not long ago we presented an analysis of the transversity distribution functions using models of hadron structure and Leading Order evolution. In the meantime Next to Leading Order evolution equations for h_1 are also available. We here present a comparison between the LO and the NLO results which confirms our previous conclusions.

Pacs: 12.39-x, 13.60.Hb, 13.88+e

Keywords: hadrons, partons, transversity, evolution.

scopetta@titan.ific.uv.es

vicente.vento@uv.es

[†]Supported in part by DGICYT-PB94-0080, DGICYT-PB95-0134 and TMR programme of the European Commisison ERB FMRX-CT96-008

In recent work we studied the transversity distributions using models of hadron structure [1]. Our scheme consists of two well defined procedures. Firstly we calculate the leading twist contributions in a low energy model of hadron structure. These results are associated with a low Q^2 , the so called hadronic scale [2]. Secondly we move our results from the hadronic scale to the experimental conditions using perturbative QCD evolution [3]. In our previous work we used Leading Order evolution, since this was the only one available for the transversity distributions at that time. Within a short period of time three calculations have appeared which allow the generalization of the evolution procedure to Next to Leading Order in this case [4, 5, 6]. It can be shown that the last two calculations agree with each other but not with the first one. Therefore we shall assume that the correct result is that of refs.[5, 6] and proceed to analyze if the conclusions of our paper are maintained to this order.

We discuss the results for the proton in two models

- i) The non relativistic model of Isgur-Karl [7];
- ii) The relativistic MIT bag model [8].

In both cases we will use the corresponding support correction as defined in [3] and [2], respectively. In Fig. 1 we show the results corresponding to g_1 and h_1 for the IK model, corresponding to the pure valence quark hadronic scenario of ref.[3], characterized by a very low hadronic scale¹. The initial data are evolved to 10 GeV², by using in the Next to Leading evolution the $\overline{MS}$ factorization scheme of refs. [5, 6] and the evolution parameters of ref. [3].

The figure shows that the diverse evolution properties of these two structure functions lead to a large difference between the two initially identical functions. The difference occurs at small x and has been noted also by other authors [9, 10]. We compare in this case the LO and NLO evolutions of g_1 and h_1 and conclude that the results are quite stable with respect to the perturbative evolution. Similar results hold for the other model.

We next turn another important result of our previous paper, namely the analysis of Soffer's inequalities. As has been pointed out by Vogelsang [5], in NLO the factorization scheme plays an important role in defining them. All the results shown will be in the $\overline{MS}$ scheme.

Both models verify the primitive Soffer inequality [11], i.e.,

$$2|h_1^q(x, Q^2)| \leq g_1^q(x, Q^2) + f_1^q(x, Q^2), \quad (1)$$

not only at the hadronic scale, but also as we evolve the distribution function towards the experimental regime. However Soffer argued [11] that his positivity bound could be used combined with data to limit the validity of models. In particular by imposing the simple relation

$$\Delta u(x) = u(x) - d(x) \quad (2)$$

¹As discussed in ref [3], the hadronic scale for LO is $\mu_0^2 = 0.079$ GeV², while for the NLO case is $\mu_0^2 = 0.094$ GeV², parameters which are fixed by assuming that the valence quarks carry all the momentum.

proposed in [12] and which is well supported by the data, it is possible to use the positivity bound to obtain the allowed range of values for h_1^u , namely

$$u(x) - d(x) \geq |h_1^u(x)| . \quad (3)$$

The MIT bag model fails the bound for large values of x [11]. We show in Fig. 2 the comparison of the experimental constraint at 4 GeV² with the Isgur-Karl and MIT bag calculations. In the figure the allowed region is described by taking the lefthand side of eq. (3) from the data. The remaining curves represent the righthand side of the equation which we have calculated from the models. It is clear from the figure that, at the hadronic scale, neither fulfils the constraint. However the above inequalities are valid at the scale of the data, then we must compare only after evolving the model calculations at NLO order from the hadronic scale to that of the data (4 GeV²). As the figure shows, consistency is achieved after this procedure. This result does not imply that the conventional models of hadron structure taken as a description of the physics at the hadronic scale are quantitatively succesful in explaining the deep inelastic data. As stated in previous analysis [3], these models give a qualitative description, which we have confirmed for the Soffer inequalities. However in order to obtain a quantitative description additional ingredients have to be added (see [1, 13] for a further discussion).

The use of models of hadron structure to describe the deep inelastic properties of the proton and neutron has proven successful for the chiral-even twist two structure functions [3] (and references therein). Several authors have generalized the analysis to the transversity functions [9, 14, 15, 16]. Since these have not been measured, this analysis has the added value of prediction. We have completed the spectrum of possible calculations by including that of a well established not relativistic model, with a fine tuned technique for constructing the structure functions and performing the RGE evolution [1]. Moreover we have returned to the highly succesful field theoretic approach of the MIT bag model and reanalyzed some of the features questioning its validity, i.e., the highly discussed Soffer inequalities [1]. In this note we have simply checked the stability of our results by performing evolution to the Next to Leading Order and therefore we have corroborated the conclusions of our previous work.

Acknowledgements

We have mantained iluminating discussions with Marco Traini and acknowledge some tuition on the evolution code. One of us (S.S.) thanks A. Drago, R. Jacob and P.J. Mulders for fruitful discussions.

References

- [1] S. Scopetta and V. Vento, FTUV-29, IFIC97-29, hep-ph/9706413.

- [2] R.L. Jaffe and G.G. Ross, Phys. Lett. B 93 (1980) 313.
- [3] M. Ropele, M. Traini and V. Vento, Nucl. Phys. A 584 (1995) 634; M. Traini, V. Vento, A. Mair and A. Zambarda, Nucl. Phys. A 614 (1997) 472.
- [4] S. Kumano and M. Miyama, hep-ph/9706420.
- [5] W. Vogelsang, hep-ph/9706511.
- [6] A. Hayashigaki, Y. Kanazawa and Yuji Koike, hep-ph/9707208.
- [7] N. Isgur and G. Karl, Phys. Rev. D 18 (1978) 4187, D 19 (1979) 2653, D 23 (1981) 817(E).
- [8] T. DeGrand, R.L. Jaffe, K. Johnson and J. Kiskis, Phys. Rev. D 12 (1975) 2060.
- [9] V. Barone, T. Calarco and A. Drago, Phys. Lett. B 390 (1997) 287.
- [10] V. Barone, University of Turin preprint DFTT-68-96, hep-ph 9703343.
- [11] J. Soffer, Phys. Rev. Lett. 74 (1995) 1292.
- [12] C. Bourrely and J. Soffer, Phys. Rev. D 51 (1995) 2108.
- [13] S. Scopetta and V. Vento, in preparation.
- [14] M. Stratmann, Z. Phys. C 60 (1993) 763.
- [15] R.L. Jaffe and X. Ji, Phys. Rev. D 49 (1994) 114.
- [16] R. Jacob, P.J. Mulders and J. Rodrigues, NIKHEF 97-018, VUTH 97-7, hep-ph/9704335.

Captions

Figure 1: We show the transversity function $h_1(x, \mu_0^2)$ (continuous line) which coincides with the spin distribution function $g_1(x, \mu_0^2)$ for the Isgur-Karl model [7] at the hadronic scale; their evolved (LO and NLO) distributions $h_1(x, Q^2)$ (dotted and long-dashed) and $g_1(x, Q^2)$ (dot-dashed and dashed) at $Q^2 = 10 \text{ GeV}^2$ are shown. The (NLO) evolutions have been performed in the $\overline{MS}$ factorization scheme [5, 6] and using the parameters taken from ref. [3] (cf. Fig. 1 (a) in ref. [1]).

Figure 2: The allowed region of Soffer determined from the experimental data at 4 GeV^2 , corresponds to the region inside the continuous line. Fig. a): the dashed line corresponds to the pure Isgur-Karl model calculation [7]; the dot-dashed line represents the evolved IK solution from an hadronic scale of 0.094 GeV^2 , while the dotted line corresponds to the evolved solution of the IK model supplemented by gluons from an hadronic scale of 0.3 GeV^2 . Fig b): the dashed line corresponds to the pure MIT calculation; the dotted line assumes a hadronic scale of 0.75 GeV^2 , while the dot-dashed line one of 0.094 GeV^2 . Stratmann analysis [14]. The evolution has been carried out to Next to Leading Order in the $\overline{MS}$ factorization scheme and the evolution parameters have been taken from ref. [3] (cf. Fig. 4 in ref. [1]).




FIGURE 1




FIGURE 2